

2.2 Study site

The study conducted using old information retroactively to malaria disease for the period of 2005 to 2008. All the data concerned with the condition have obtained from the Ministry of Health of South Darfur state, from malaria and neglected tropical diseases program. As the population number received from the Department of Statistics in South Darfur state, the 2008 census and the backwards projections for the following years until 2005. Maps have varied from department of surveying in South Darfur state. The data classified, processed and analyzed by the use of GIS tool, i.e., ArcGIS (V10.1; ESRI). After the appropriate data equipping. The data processed and analyzed using the Geographic Information System GIS to produce digital maps. These maps are: ABIR, IR, FR, and Pf%.

3 Results

The spatial distribution changes for seven years of malaria in three localities in Kass zone are illustrated by digital maps according to two parameters; ABIR, IR, FR, and PF.

Figure 2 indicates the changes of ABIR in 3 localities of Kass zone, South Darfur Sudan from 2005 to 2008. ABER takes into account the efficiency and adequacy of case detection mechanisms [9]. Within a 4 year periods, values of ABER in Kass locality were constantly reported >10% through the whole study period. The changes of malaria in IR for three localities in Kass zone from 2005 to 2008 graphically shown in Figure 3. Within 4 years duration, Kass locality reported IR >10/10000 population in 2005 and 2008. Sharq Gabal reported <10/10000 population from 2005-2007. Meanwhile, Shattai locality reported IR >10/10000 in 2006 and 2007.

Figure 4 shows the changes in FR on 3 localities in Kass zone from 2005 to 2008. There was malaria fatality cases reported in Kass locality FR > 10% during the study period. Figure 5 indicates the changes in Pf for 4 localities in Kass zone from 2005 to 2008. There was malaria Pf cases reported 71-100% in Kass locality, but Sharq Gabal and Shattai reported Pf <30% in two years for each through the study period.




Fig 2: ABER from 2005-2008 in Kass zone

Fig 3: IR from 2005-2008 in Kass zone




Fig 4: FR from 2005-2008 in Kass zone

Fig 5: Pf% from 2005-2008 in Kass zone

3. Discussions

In this study, the results have demonstrated a large measurements data set among 14832 patients in Kass zone, about 32% <5 years old and 68% >5 years old. The blood checked 24% out of the whole patients. The infections are about 42% out of the checked patients. ABER measures coverage of the surveillance program, and potentially local fever incidence. The distribution of ABER through the four years from (2005 – 2008) in Kass zone indicates that the study area is highly endemic. The higher prevalence of malaria found in Kass locality, IR, ABIR, and Pf was recorded through the four FR study years, which is the highest case detection in the zone. Because Kass locality is the compound of the camps for displaced persons due to tribal and armed militia groups. Through the entire period, we can see the decreasing of IR towards 2008. Because of the policy of indoor residual spraying (IRS) to sensitive areas and the distribution of insecticide-treated nets (ITNs) in most of Kass localities. However, IR still very high and needs more treating and investigations. Relating to the FR, the malaria fatality cases reported in all Kass zone. Reasons and factors that helps a lot in the presence of death cases that emerged during the study period are, Some delay of the patients visiting the clinics, especially in rural areas, some laboratories failed in the diagnosis of the quality and stage of the disease, leading to failure to determine the appropriate treatment on time. As well as natural factors, such as torrential rain that was hindering malaria control teams and deliver medicines, moreover the lack of paved roads to be provided. There is also a dearth of means of storage of medications on sweeteners for lack of electricity stability [10,11].

Regarding all of the above consequences, we find that the malaria control program in Kass zone is active to some extent, but it still needs more studies and intensive programs. To combat malaria, the government must implement several strategies, such as early case detection, close monitoring of important cases, using bed nets, monitoring of drug resistance, residual spraying, and collaborating with the neighboring states and countries.

4. Conclusions

It clearly appears the change in the spatial distribution of malaria within a 4-year periods, regarding the digital mapping of infections in South Darfur State.

The findings indicate the irregularity variation in malaria distribution through the study period, as well as in the diverse malaria localities. Mainly malaria disease is brought up in the central part of the of Kass zone especially in Kass locality. Our conclusion notes that Kass zone still an endemic area and need more studies and a comprehensive program to decrease the malaria infection.

5. Acknowledgement

The authors would like to acknowledge the Malaria and Neglected Tropical Diseases, Ministry of Health, South Darfur, Sudan, Awadalla Musa, and Ibrahim Adam.

6. Authors' contributions

HEH conceived, designed the study, performed the data collection, and prepared the final draft. YB supervised the data analysis, reviewed the manuscript critically, and contributed intellectual input. All authors read and approved the final manuscript.

7. Conflicts of interest

The authors declare that they do not have competing interests.

8. References

- [1] Organization WH. Antimicrobial resistance global report on surveillance: 2014 summary. 2014.
- [2] Malik E, Atta HY, Weis M, et al. Sudan Roll Back malaria consultative mission: essential actions
- [3] Abdalla SI, Malik EM, Ali KM. The burden of malaria in Sudan: incidence, mortality and disability–adjusted life–years. *Malaria journal* 2007, 6(1):1.
- [4] Adam I, Elhassan EM, Haggaz AED, et al. A perspective of the epidemiology of malaria and anaemia and their impact on maternal and perinatal outcomes in Sudan. *The Journal of Infection in Developing Countries* 2010, 5(02):83-87.
- [5] Awad M, Alkadru A, Behrens R, et al. Descriptive study on the efficacy and safety of artesunate suppository in combination with other antimalarial in the treatment of severe malaria in Sudan. *The American journal of tropical medicine and hygiene* 2003, 68(2):153-158.
- [6] Bi Y, Hu W, Liu H, et al. Can slide positivity rates predict malaria transmission? *Malaria journal* 2012, 11(1):1.
- [7] Hassan HE. *Malaria Control Using GIS (Khartoum State)*. Sudan University of Science &Technology; 2008.
- [8] Musa MI, Shohaimi S, Hashim NR, et al. A climate distribution model of malaria transmission in Sudan. *Geospatial health* 2012, 7(1):27-36.
- [9] Malaria and VHF Outbreak in Darfur, Sudan, Situation Report No 14: 18 January - 7 February 2016 Federal Ministry of Health | World Health Organization. <https://reliefweb.int/report/sudan/malaria-and-vhf-outbreak-darfur-sudan-situation-report-no-14-18-january-7-february-2016>
- [10] Malik EM, Khalafalla O. Malaria in Sudan: past, present and the future. *Gezira journal of health sciences* 2004, 1(1 Sup).

- [11] Nmor JC, Sunahara T, Goto K, et al. Topographic models for predicting malaria vector breeding habitats: potential tools for vector control managers. *Parasites & vectors* 2013, 6(1):1.

IEEESEM