

indolizino [1,2-c]quinolin- 7(5H) mono-[4(1)]-zero-13[4(1)]methine (**7a -f**) are highly coloured compounds ranging from brownish-red / red, dark brown / deep violet (blue). They are partially soluble in non-polar solvents and easily soluble in polar solvents exhibiting highly coloured solutions with slight or intense green fluorescence depending upon the incorporating anhydro base of N-bridge head heterobicyclic moieties or heterocyclic quaternary residue. Compounds (**4,5a-d, 6a-f & 7a -f**) are highly soluble in concentrated sulphuric acid liberating iodine vapour on warming. Their ethanolic solution turned yellow or became colourless on acidification, while their permanent colour was restored on basification. The absorption spectra of benzo[g]indolizino[1,2-c]quinolin-7(5H)mono-[4(1)]-zero-13[4(1)]methine (**4a-d**) in 95% ethanol showed absorption bands batho- chromically (hypsochromically) shifted depending upon the nature of heterocyclic quaternary residue **A(B)**, their linkage position. Thus, the absorption spectra of dye [**4a**, A (B) = pyridin-4-ium (quinolin-1-ium)] showed λ_{\max} 455,370 nm (ϵ_{\max} , 2000,4000 mol⁻¹cm²). Substituting of **A(B)** = pyridin-4-ium(quinolin-1-ium) in dye (**4a**) by **A (B)**= quinolin-4-ium(quinolin-1-ium) in dye (**4b**) resulted in bathochromically shifted by (5-10 nm) than (**4a**) [**4b**, λ_{\max} 465, 375 nm, ϵ_{\max} , 8000,3400 mol⁻¹cm²]. This is due to the more extensive π -delocalization within benzene ring attached to pyridine ring in dye (**4b**). Substituting of **A (B)** = quinolin-4-ium (quinolin-1-ium) in dye (**4b**) by **A (B)** = quinolin-1-ium (quinolin-1-ium) in dye (**4c**) causes hypso-chromically shifted of 5nm than (**4b**) [**4c**, λ_{\max} 460,373 nm ϵ_{\max} , 4000,7400 mol⁻¹ cm²]. This is due to an easier charge transfer within extended π -delocalization of 4-ium salt rather than 1-ium **Table (2)**. Substituting **A (B)** =quinolin-4-ium (quinolin-1-ium) in dye (**4b**) by A (B) =quinolin-4-ium (pyridin-1-ium) in dye (**4d**) move the absorption bands towards batho- chromic shift by10 nm than (**4b**). [**4d**, λ_{\max} 475 nm ϵ_{\max} , 13200 mol⁻¹ cm²]. This is due to that inserting of ring causes the more extensive π -delocalization and extra conjugation, so the absorption takes place at longer wavelength, **Table (2)**. The absorption spectra of benzo[g]indolizino[1,2-c]quinolin-7(5H)mono-[4(1)]-zero-13[4(1)]methine (**5a-d**) in 95% ethanol showed absorption bands batho- chromically (hypsochromically) shifted depending upon the nature of heterocyclic quaternary residue **A(B)**, their linkage position. Thus, the absorption spectra of dye [**5a**, A (B) =pyridin-4-ium (quinolin-1-ium)] showed λ_{\max} 510,445,415 nm (ϵ_{\max} , 6000, 8000, 9600 mol⁻¹ cm²). Substituting of **A (B)** = pyridin-4-ium (quinolin-1-ium) in dye (**5a**) by **A (B)** =quinolin-4-ium(quinolin-

1-ium) in dye **(5b)** resulted in bathochromically shifted by (80nm) than **(5a)** [**5b**, λ_{\max} 660, 570, 470, 435 nm, ϵ_{\max} , 2400,5400,7800,8800 mol⁻¹cm²]. This is due to the more extensive π -delocalization within benzene ring attached to pyridine ring in dye **(5b)**. Substituting of **A (B)** =quinolin-4-ium (quinolin-1-ium) in dye **(5b)** by **A (B)** = quinolin-1-ium (quinolin-1-ium) in dye **(5c)** causes hypsochromically shifted of 45nm than **(5b)** [**5c**, λ_{\max} 512,450 nm ϵ_{\max} , 6800,14000 mol⁻¹ cm²]. This is due to an easier charge transfer within extended π -delocalization of 4-ium salt rather than 1-ium


Table (3). Substituting **A (B)** =quinolin-4-ium (quinolin-1-ium) in dye **(5b)** by **A (B)** =quinolin-4-ium (pyridin-1-ium) in dye **(5d)** move the absorption bands towards bathochromic shift by10nm than **(5b)**. [**5d**, λ_{\max} 690, 595, 560,400nm ϵ_{\max} , 3200,12800,11400,9600 mol⁻¹ cm²]. This is due to that inserting of ring causes the more extensive π -delocalization and extra conjugation, so the absorption takes place at longer wavelength, **Table (2)**. The absorption spectra of benzo [g] indolizino[1,2-c]quinolin-7(5H)mono-[4(1)]-zero-13[4(1)]methine **(6a-f)** in 95% ethanol showed absorption bands bathochromically (hypsochromically) shifted depending upon the nature of heterocyclic quaternary residue **A[A` (B)]**, their linkage position. Thus, the absorption spectra of dye **[6a, A [A` (B)]** pyridin-4-ium [quinolin-4-ium (quinolin-1-ium)] showed λ_{\max} 610, 380 nm (ϵ_{\max} , 1800,9800 mol⁻¹ cm²). Substituting of **A[A` (B)]**pyridin-4-ium[quinolin-4-ium (quinolin-1-ium)] in dye **(6a)** by **A[A` (B)]**quinolin-4-ium[quinolin-4-ium (quinolin-1-ium)] in dye **(6b)** resulted in bathochromically shifted by (80nm) than **(6a)** [**6b**, λ_{\max} 690,575,460, ϵ_{\max} , 2900,6000,4200 mol⁻¹cm²]. This is due to the more extensive π -delocalization within benzene ring attached to pyridine ring in dye **(6b)**. Substituting of **A [A` (B)]** quinolin-4-ium[quinolin-4-ium(quinolin-1-ium)] in dye **(6b)** by **A[A` (B)]** quinolin-1-ium[quinolin-4-ium, (quinolin-1-ium)] in dye **(6c)** causes hypsochromically shifted of 45nm than **(6b)** [**6c**, λ_{\max} 635sh,465,390 nm ϵ_{\max} , 2500,2000, 7000 mol⁻¹ cm²]. This is due to an easier charge transfer within extended π -delocalization of 4-ium salt rather than 1-ium

Table (3). Substituting **A [A` (B)]** quinolin-4-ium [quinolin-4-ium (quinolin-1-ium)] in dye **(6b)** by **A [A` (B)]**quinolin-4-ium[quinolin-4-ium(pyridin-1-ium)] in dye **(6d)** move the absorption bands towards bathochromic shift by10nm than **(6b)**. [**6d**, λ_{\max} 700, 600, 470 nm ϵ_{\max} , 2600,4600,10800 mol⁻¹ cm²]. This is due to that inserting of ring causes the more extensive π -delocalization and extra conjugation, so the absorption takes place at longer wavelength, **Table (3)**. The absorption spectra of **(6e,f)** in 95%

ethanol showed absorption bands at **[6e, A [A` (B)] quinolin-4-ium, [pyridin-4-ium (quinolin-1-ium)] (6e, λ_{\max} 470,335 nm, ϵ_{\max} , 3500, 5500 mol⁻¹ cm²]**, **[6f, A[A` (B)] quinolin-4-ium [quinolin-1-ium (quinolin-1-ium)] [6f, λ_{\max} 500 nm, ϵ_{\max} , 13000 mol⁻¹ cm²]**. It was obvious that the absorption spectra of **(6f)** showed absorption bands bathochromically shifted by (30nm) than **(6e)** and absorb in the green-blue light region. This is due to the extensive of π -delocalization and an increasing in conjugation within quinolin-1-ium resulted in a charge transfer from ring junction N-bridge atom towards N-heterocyclic quaternary residue. The absorption spectra of benzo[g]indolizino[1,2-c]quinolin-7(5H)mono-[4(1)]-zero-13[4(1)] methine **(7a-f)** in 95% ethanol showed absorption bands bathochromically (hypsochromically) shifted depending upon the nature of heterocyclic quaternary residue **A[A` (B)]**, their linkage position. Thus, the absorption spectra of dye **[7a, A [A` (B)] pyridin-4-ium [quinolin-4-ium (quinolin-1-ium)]** showed λ_{\max} 620, 390nm (ϵ_{\max} , 1800, 1000 mol⁻¹ cm²). Substituting of **A[A` (B)] pyridin-4-ium[quinolin-4-ium(quinolin-1-ium)]** in dye **(7a)** by **A [A` (B)] quinolin-4-ium[quinolin-4-ium(quinolin-1-ium)]** in dye **(7b)** resulted in bathochromically shifted by (70nm) than **(7a)** **[7b, λ_{\max} 690, 585, 470, ϵ_{\max} , (36000), (50000) (50000) mol⁻¹ cm²]**. This is due to the more extensive π -delocalization within benzene ring attached to pyridine ring in dye **(7b)**. Substituting of A [A` (B)] quinolin-4-ium[quinolin-4-ium(quinolin-1-ium)] in dye **(7b)** by A[A` (B)] quinolin-1-ium[quinolin-4-ium, (quinolin-1-ium)] in dye **(7c)** causes hypsochromically shifted of 45nm than **(7b)** **[7c, λ_{\max} 645sh, 475, 400 nm ϵ_{\max} , 7000 ,13500 , 7100 mol⁻¹ cm²]**. This is due to an easier charge transfer within extended π -delocalization of 4-ium salt rather than 1-ium **Table (3)**. Substituting A [A` (B)] quinolin-4-ium [quinolin-4-ium(quinolin-1-ium)] in dye **(7b)** by A [A` (B)]quinolin-4-ium, [quinolin-4-ium(pyridin-1-ium)] in dye **(7d)** move the absorption bands towards bathochromic shift by10nm than **(7b)**. **[7d, λ_{\max} 700, 620,490 nm ϵ_{\max} , (5000, 7500, 10800 mol⁻¹ cm²]**. This is due to that inserting of ring causes the more extensive π -delocalization and extra conjugation, so the absorption takes place at longer wavelength, **Table (3)**. The absorption spectra of **(7e,f)** in 95% ethanol showed absorption bands at **[7e, A [A` (B)] quinolin-4-ium, [pyridin-4-ium (quinolin-1-ium)] (λ_{\max} 480, 355 nm, ϵ_{\max} , 3500, 5000 mol⁻¹ cm²]**, **[7f, A[A` (B)] quinolin-4-ium [quinolin-1-ium (quinolin-1-ium)] λ_{\max} 520 nm, ϵ_{\max} , 5500 mol⁻¹ cm²]**. It was obvious that the absorption spectra of **(7f)** showed absorption bands bathochromically shifted by (40nm) than **(7e)** and absorb in the green-blue light

region. This is due to the extensive of π -delocalization and an increasing in conjugation within quinolin-1-ium resulted in a charge transfer from ring junction N-bridge atom towards N-heterocyclic quaternary residue. On comparison of the absorption spectra of dyes (**6d**, **7d**), it was obvious that the absorption bands of [**7b**, λ_{\max} 700, 620, 490 nm ϵ_{\max} , (5000, 7500, 10800 mol⁻¹ cm²)] move to bathochromic shift of 20nm than those of [**6d**, λ_{\max} 700, 600, 470 nm ϵ_{\max} , 2600, 4600, 10800 mol⁻¹ cm²], **Table (3)**. This is due to that the ring junction nitrogen atom in the former dye affected on the charge transfer by both electron releasing inductive and mesomeric effects better than those of the latter dye in which the ring junction nitrogen atom represents as electron withdrawing character (electron withdrawing inductive effect) due to its neighboring. Additionally, the former dye incorporates an endocyclic quinoid structure causes an increasing in π -delocalization better than the latter dye incorporating an exocyclic quinoid analogous. **Table (3)**. On comparison of the absorption spectra of dyes (**5a-d**, **7a-f**), it was obvious, that the absorption spectra of (**7e,f**) in 95% ethanol showed absorption bands in of great values λ_{\max} and ϵ_{\max} than those of (**5a-d**). This is due to of the more multi internal charge transfer pathways of (**7a-f**) than those of (**5a-d**), as was observed in **Equation (2)**, at [**7e**, A [A` (B)] quinolin-4-ium, [pyridin-4-ium (quinolin-1-ium)] (λ_{\max} 480, 355 nm, ϵ_{\max} , 3500, 5000 mol⁻¹ cm²), [**7f**, A[A` (B)] quinolin-4-ium [quinolin-1-ium (quinolin-1-ium)] λ_{\max} 520 nm, ϵ_{\max} , 5500 mol⁻¹ cm²]. It was obvious that the absorption spectra of (**7f**) showed absorption bands bathochromically shifted by (40nm) than (**7e**) and absorb in the green-blue light region. This is due to the extensive of π -delocalization and an increasing in conjugation within quinolin-1-ium resulted in a charge transfer from ring junction N-bridge atom towards N-heterocyclic quaternary residue. Meanwhile, the absorption spectra of dye [**5a**, A (B) =pyridin-4-ium (quinolin-1-ium) showed λ_{\max} 510, 445, 415 nm (ϵ_{\max} , 6000, 8000, 9600 mol⁻¹ cm²). Substituting of **A (B)** = pyridin-4-ium (quinolin-1-ium) in dye (**5a**) by **A (B)** =quinolin-4-ium(quinolin-1-ium) in dye (**5b**) resulted in bathochromically shifted by (80nm) than (**5a**) [**5b**, λ_{\max} 660, 570, 470, 435 nm, ϵ_{\max} , 2400, 5400, 7800, 8800 mol⁻¹ cm²]. This is due to the more extensive π -delocalization within benzene ring attached to pyridine ring in dye (**5b**). Substituting of **A (B)** =quinolin-4-ium (quinolin-1-ium) in dye (**5b**) by **A (B)** = quinolin-1-ium (quinolin-1-ium) in dye (**5c**) causes hypso- chromically shifted of 45nm than (**5b**) [**5c**, λ_{\max} 512, 450 nm ϵ_{\max} , 6800, 14000 mol⁻¹ cm²]. This is due to an easier charge

transfer within extended π -delocalization of 4-ium salt rather than 1-ium **Table (3)**. Substituting **A (B)** =quinolin-4-ium (quinolin-1-ium) in dye **(5b)** by A (B) =quinolin-4-ium (pyridin-1-ium) in dye **(5d)** move the absorption bands towards bathochromic shift by 10nm than **(5b)**. [**5d**, λ_{\max} 690, 595, 560, 400nm ϵ_{\max} , 3200, 12800, 11400, 9600 mol⁻¹ cm²]. This is due to that inserting of ring causes the more extensive π -delocalization and extra conjugation, so the absorption takes place at longer wavelength, **Table (4)**.



Equation (2): Internal Charge Transfer Pathways of (5a-d and 7a-f)

The high absorption spectra of **(5a-d)** value range λ_{\max} n.m 510,445,415, 660,570,470,435,512, 450, 690,595,560,400 ϵ_{\max} (6000), (8000), (9600), (2400), (5400),(7800),(8800),(6800),(14000),(3200),(12800),(11400),(9600) mol⁻¹ cm² **Table (3)**, is due to the interpretation of multi [ICT] pathways of **(7a-d)** is due to the different orientation of the internal charge transfer for more directions, it is from the direction of multi [ICT] internal charges is from NH of piperidine nuclei as electron source towards heterocyclic quaternary salts as electron sink of zero methine cyanine type, as was represents with blue arrows colour or from NH of piperidine nuclei as electron source towards either naphtho quinone cyclic carbonyl, as was represents with green arrows colour or towards Benz indolizinium iodide, as was represents with red arrows colour as electron sink.

Meanwhile, the high absorption spectra of **(7a-f)** value range at λ_{\max} nm 390-620, 470-690, 400-645, 490-700, 355-480, 520, ϵ_{\max} (1000, 36000, 50000, 50000, 7000, 13500, 5000, 7500, 3500, 5000, 5500, 7900, 7900) $\text{mol}^{-1} \text{cm}^2$ **Table (3)**, is due to the interpretation of multi **[ICT]** of **(7a-d)** is due to the different orientation of the internal charge transfer for more directions, it is from the N-bridge head of indolizine nuclei as electron source either towards cyclic carbonyl group of quinoid as electron sink, as was represents with blue arrows colour, or towards heterocyclic quaternary salts as electron sink of mono methine cyanine type, as was represents with black arrows colour. The other direction of multi **[ICT]** internal charges is from NH of piperidine nuclei as electron source towards heterocyclic quaternary salts as electron sink of zero methine cyanine type, as was represents with red arrows colour or towards heterocyclic quaternary salts as electron sink of mono methine cyanine type, as was represents with green arrows colour. The calculations and computational studies including **HOMO-LOMO** of **(5a-d & 7a-f)** were under investigation to elucidate such activities by one of my colleagues in future.

3. EXPERIMENTAL

All melting points are uncorrected. Elemental analysis were carried out at the micro analytical centre (Cairo-university). The IR spectra were determined with perkin elmer infrared 127B spectrophotometer (Cairo– University). The $^1\text{H-NMR}$ spectra were recorded with EM-390 90 MHz NMR spectrophotometer. (Cairo– University). Mass spectra were recorded on a Hp Ms 6488 spectrometer. The visible were recorded on UV-visible recording spectrophotometer UV-240 and UV-160 A. Faculty of Science Aswan 4-(2-(3-amino-1, 4-di-oxo-1,4-di[H] naphthalen-2-yl)-2-oxo-ethyl)-1-ethyl-pyridin[quinolin-4(1)]-1-ium iodide **(1a-c)** was prepared in a way that described in prospective reference [12].

3.1. Synthesis of 1-(2-(3-amino-1, 4-dioxo-1, 4-di [H]naphthalen-2-yl)-1-(1-ethyl-pyridin [quinolin-4(1)]-1-ium)-2-oxo-ethyl)-2-methyl-pyridin (quinolin)-1-ium iodide, (2a-c).

An ethanolic solution of **(1a, b,** and 0.01mol) and 2-methyl-pyridin [quinoline]-bases (0.01mol) in iodine was refluxed for 3-5 hrs. The reaction mixtures were filtrated from unreacted materials. The filtrate concentrated to one third of its volume, cooled and acidified with acetic acid. The precipitated products after dilution with water were separated, filtrated, crystallized from ethanol to give **(2a-c), Table (1).**

IR (KBr cm^{-1}) of **(2b)** showed general absorption peaks at 2890 cm^{-1} (ylide anion), $2921\text{-}2933 \text{ cm}^{-1}$ (γ H-Q-salt), $3150\text{-}3000 \text{ cm}^{-1}$ (γ C-H, γ N-methyl-pyridinium iodide), $1715\text{-}1730 \text{ cm}^{-1}$ (γ acyclic C=O), 1600 cm^{-1} (γ C=C conj. Cyclic), $3350\text{-}3310 \text{ cm}^{-1}$, (γ N-H, stretch primary amine), and (γ N-H str., pyridinium ion), $1631\text{-}1625 \text{ cm}^{-1}$ (γ quaternary pyridine), $1631\text{-}1625 \text{ cm}^{-1}$. (γ pyridinium salts), $1129\text{-}092 \text{ cm}^{-1}$ (γ exocyclic N-C str. pyridinium salts), 1715 cm^{-1} (γ quinoid cyclic C=O, $3150\text{-}3000 \text{ cm}^{-1}$ (γ C-H. γ N-ethyl-pyridinium iodide), $3037\text{-}3144 \text{ cm}^{-1}$ (γ pyridinium quaternary), $1129\text{-}092 \text{ cm}^{-1}$ (γ exocyclic N-C str. pyridinium salts), $3037\text{-}3144 \text{ cm}^{-1}$ (quaternary salts), $1129\text{-}092 \text{ cm}^{-1}$ (γ exocyclic N-C-alkyl str. pyridinium salts), Characteristic strong $1631\text{-}625 \text{ cm}^{-1}$ (γ C=N of heterocyclic quaternary nitrogen atom or ring vibrations of pyridinium ion) [13]... $^1\text{H-NMR}$ (DMSO, 300 MHz) of selected **(2b)** showed in addition to the general signals at δ 7.60-7.72 ppm (m, 4H, Ar-H), δ 7.67-8.90 ppm (m, 6H, quinolin-4-ium), δ 8.10-9.11 ppm (m, 6H, quinolin-1-ium), δ 1.57 ppm (t, 3H, CH_3), δ 4.80 ppm (q, 2H, CH_2), characteristic signal at δ 3.6 ppm (s, 1H, CH), δ 2.92 ppm (s, 3H, CH_3) & δ 4.79 ppm (s, 2H, NH_2) [14,15].. Mass spectrum of **(2b)** reveals a molecular ion, peaks at $m/z = 766$ and base peak at $m/z = 200$ [16].

3.2. Synthesis of 1-(2-(1,4-dioxo-3-(prop-1-en-2-ylamino)-1,4-dihydro)naphthalen-2-yl)-1-(1-ethyl-pyridin[quinolin-4(1)]-1-ium-4-yl)-2-oxo-ethyl)-2-(2-oxo-propyl)pyridin (quinolin)-1-ium, (3a-d)

A mixture of **(2a-d, 0.005 mol)** and acetic anhydride (10 ml) was refluxed for 3 hours. The reaction mixture was filtrated from unreacted materials. The filtrate was concentrated and cooled concentrated to one third of its volume, cooled and precipitated by adding of ice-water mixture to give **(3a-d), Table (1)**. IR (KBr cm^{-1}) of **(3b)** showed in addition to general absorption peaks cited for **(2b)**, characteristic absorption peaks at 1712 cm^{-1} (γ COCH_3), 1640 cm^{-1} (γ str. enolized acetyl carbonyl group), Characteristic strong $1631\text{-}625 \text{ cm}^{-1}$ (γ C=N of heterocyclic quaternary nitrogen atom or ring vibrations of pyridinium ion), 3120 cm^{-1} , (γ N-H str., pyridinium ion), $1631\text{-}1625 \text{ cm}^{-1}$ (γ quatern pyridine), $1631\text{-}1625 \text{ cm}^{-1}$. (γ pyridinium salts), $1129\text{-}092 \text{ cm}^{-1}$ (γ exocyclic N-C str. pyridin-ium salts), 1715 cm^{-1} (γ quinoid cyclic C=O, $3150\text{-}3000 \text{ cm}^{-1}$ (γ N-ethyl-pyridin-ium iodide), $3037\text{-}3144 \text{ cm}^{-1}$ (γ 2 bands pyridinium quaternary), $1129\text{-}092 \text{ cm}^{-1}$ (γ exocyclic N-C str. pyridinium salts), $3037\text{-}3144 \text{ cm}^{-1}$ (quaternary salts), $1129\text{-}092 \text{ cm}^{-1}$ (γ exocyclic N-C-alkyl str. pyridinium salts) [13]. $^1\text{H-NMR}$ (DMSO, 300 MHz) of selected **(3b)** showed in addition to the general signals at δ 7.60-7.72 ppm (m, 4H, Ar-H), δ 7.67-8.90 ppm (m, 6H, quinolin-4-ium),

δ 8.10-9.11 ppm (m, 6H, quinolin-1-ium), δ 1.57 ppm (t, 3H, CH₃), δ 4.80 ppm (q, 2H, CH₂), characteristic signal at δ 3.6 ppm (s, 1H, CH), δ 1.86-1.89 ppm (s, 3H, CH₃ of acetyl group) & δ 9.30 ppm (s, 1H, NH), δ 3.71 ppm (s, 2H, CH₂) [14,15]. Mass spectrum of (3b) reveals a molecular ion at $m/z = 773$, and base peak at $m/z = 242.05$ [16].

3.3. Synthesis of Self-Assembly Heterocyclic [ICT] functional dye 6-acetyl-7-methyl-5, 14-dioxo-6, 14-di [H]-5H-benzo[g]indolizino [1, 2-c] quinolin-12-ium iodide-zero-13[4(1)] methine cyanine dyes, (4a-d),

An Ethanolic solution of (3a-d, 0.005 mol) in the presence of few mls. Piperidine was refluxed for 3 hours. The reaction mixture was filtrated from unreacted materials. The filtrate was concentrated and cooled concentrated to one third of its volume, cooled and precipitated by adding of ice-water mixture. to give (4a-d), Table (2). IR ($\nu^{\text{KBr}} \text{ cm}^{-1}$) of (4b) showed in addition to the general absorption peaks at (ν CH₃ Stretching Vibration), at 2890 cm^{-1} (ν ylide anion), 3150-3000 cm^{-1} (ν N-ethyl-pyridin-ium iodide), 1715 cm^{-1} (ν quinoid cyclic C=O), groups, characteristic absorption peaks at 1712 cm^{-1} (ν (NCOCH₃), for (4b) & disappearance 1712 cm^{-1} (ν (NCOCH₃) [13]. ¹H-NMR (DMSO, 300 MHz) of selected (4b) showed in addition to the general signals at δ 7.60-7.72 ppm (m, 4H, Ar-H), δ 7.94-9.01 ppm (m, 6H, quinolin-4-ium), δ 7.57-9.40 ppm (m, 6H, benzo [g]indolizine), δ 1.57 ppm (t, 3H, CH₃), δ 4.80 ppm (q, 2H, CH₂), characteristic signal at δ 3.6 ppm (s, 1H, CH), δ 1.91 ppm (s, 3H, CH₃ of acetyl group) & δ 9.30 ppm (s, 1H, NH), δ 3.71 ppm (s, 2H, CH₂), δ 2.26 ppm (s, 3H, CH₃) [14,15]. Mass spectrum of (4b) reveals a molecular ion, peaks at $m/z = 818$ and base peak at $m/z = 760$ [16].

3.4. Synthesis of Self-Assembly Heterocyclic functional dye 13-(1-ethyl-pyridin-1-ium-4-yl)-7-methyl-5, 14-dioxo-6, 14-dihydro-5H-benzo[g] indolizino [1, 2-c] quinolin-12-ium iodide, (5a-d)

An ethanolic solution of (4a-d, 0.005 mol) in presence of KOH or K₂CO₃ or 1m M NaHSO₃ and use 0.1N HCl [13] was refluxed for 1 hour. The reaction mixture was filtrated from unreacted materials, concentrated and cooled; the solid product was collected and crystallized from ethanol to give (5a-d), Table (3). IR ($\nu^{\text{KBr}} \text{ cm}^{-1}$) of (5b) showed in addition to the general absorption peaks at (ν CH₃ Stretching Vibration), at 2890 cm^{-1} (ν ylide anion), 3150-3000 cm^{-1} (ν N-ethyl-pyridin-ium iodide), 1715 cm^{-1} (ν quinoid cyclic C=O), groups, characteristic appearance of absorption band at

3300-3500 cm^{-1} (ν_{NH}) for **(5b)** [13]. $^1\text{H-NMR}$ (DMSO, 300 MHz) of selected **(5b)** showed in addition to the general signals at δ 7.60-7.72 ppm (m, 4H, Ar-H), δ 7.67-8.90 ppm (m, 6H, quinolin-4-ium), δ 7.57-9.40 ppm (m, 6H, benzo[g]indolizine), δ 1.57 ppm (t, 3H, CH_3), δ 4.80 ppm (q, 2H, CH_2), characteristic signal at δ 3.6 ppm (s, 1H, CH), & δ 5.93 ppm (s, 1H, NH), δ 3.71 ppm (s, 2H, CH_2), δ 2.26 ppm (s, 3H, CH_3) [14,15]. Mass spectrum of **(5b)** reveals a molecular ion at $m/z = 773$, and base peak at $m/z = 758$ [16].

3.5. Synthesis of 4-(6-acetyl-5, 14-dioxo-6, 14-di [Hbenzo[g]indolizino [1, 2-c] quinolin-7(5H) mono-[4(1)]-zero-13[4(1)] methine, (6a-f)

An Ethanolic solution of **(4a-d)**, 0.01 mol.) and pyridin [quinolin]-4(1)-ium-ethiodide salts (0.01 moles) in piperidine was refluxed on water bath for one hour. The reaction mixture was filtrated from unreacted materials. The filtrate was concentrated and cooled. The precipitated products after dilution with water were separated, filtrated, washed with water several times and crystallized from ethanol to give **(6a-f)**, Table [4]. IR (ν^{KBr} cm^{-1}) of **(6b)** showed in addition to the general absorption peaks at (CH_3 Stretching Vibration), 2890 cm^{-1} (ν ylide anion), $3150\text{-}3000 \text{ cm}^{-1}$ (ν N-ethyl-pyridin-ium iodide), 1715 cm^{-1} (ν quinoid cyclic $\text{C}=\text{O}$), characteristic absorption peaks at 1712 cm^{-1} (ν NCOCH_3) for **(6b)** [13]. $^1\text{H-NMR}$ (DMSO, 300 MHz) of selected **(6b)** showed in addition to the general signals at δ 7.60-7.72 ppm (m, 4H, Ar-H), δ 7.67-8.90 ppm (m, 12H, quinolin-4-ium), δ 7.57-9.40 ppm (m, 6H, benzo[g]indolizine), δ 1.57 ppm (t, 3H, CH_3), δ 4.80 ppm (q, 2H, CH_2), characteristic signal at δ 3.6 ppm (s, 1H, CH), & δ 5.93 ppm (s, 1H, NH), δ 3.71 ppm (s, 2H, CH_2), δ 2.26 ppm (s, 3H, CH_3), δ 1.91 ppm (s, 3H, CH_3 of acetyl group), δ 5.99 ppm (s, 1H, =CH). $^1\text{H-NMR}$ (DMSO, 300 MHz) of selected compound **(7b)** showed in addition to the general signals at δ 7.60-7.72 ppm (m, 4H, Ar-H), δ 8.10-9.08 ppm (m, 12H, quinolin-4-ium), δ 7.77-8.48 ppm (m, 6H, benzo[g]indolizine), δ 1.57 ppm (t, 3H, CH_3), δ 4.80 ppm (q, 2H, CH_2), characteristic signal at δ 3.6 ppm (s, 1H, CH), & δ 5.93 ppm (s, 1H, NH), δ 3.71 ppm (s, 2H, CH_2), δ 2.26 ppm (s, 3H, CH_3), δ 1.91 ppm (s, 3H, CH_3 of acetyl group), δ 6.03 ppm (s, 1H, =CH) [14,15]. The structure **(6b)** was consider most likely based on mass spectrum resulted in M^+ peak at $m/e = 971$ and base peak at $m/e = 928$ which in agreement with the structure **(6b)** [16]

3.6. Synthesis of 5,14-dioxo-5,6,7,14-tetra[H]benzo[g]indolizino[1,2-c]quinolin-7(5H)mono-[4(1)]-zero-13[4(1)]methine (7a-f)

An Ethanolic solution of (**5a-d**, 0.01 mol.) and pyridin [quinolin]-4(1)-ium-ethiodide salts (0.01 moles) in piperidine was refluxed on water bath for one hour. The reaction mixture was filtrated from unreacted materials. The filtrate was concentrated and cooled. The precipitated products after dilution with water were separated, filtrated, washed with water several times and crystallized from ethanol to give (**7a-f**), **Table [4]**. IR (ν^{KBr} cm^{-1}) of (**7b**) showed in addition to the general absorption peaks at (CH_3 Stretching Vibration), 2890 cm^{-1} (ν ylide anion), $3150\text{-}3000 \text{ cm}^{-1}$ (ν N-ethyl-pyridin-ium iodide), 1715 cm^{-1} (ν quinoid cyclic C=O), characteristic absorption peaks disappearance 1712 cm^{-1} (ν (NCOCH₃) and appearance of absorption band at $3300\text{-}3500 \text{ cm}^{-1}$ (ν NH) for (**7b**)[**13**]. ¹H-NMR (DMSO, 300 MHz) of selected compound (**7b**) showed in addition to the general signals at δ 7.60-7.72 ppm (m, 4H, Ar-H), δ 8.10-9.08 ppm (m, 12H, quinolin-4-ium), δ 7.77-8.48 ppm (m, 6H, benzo[g]indolizine), δ 1.57 ppm (t, 3H, CH₃), δ 4.80 ppm (q, 2H, CH₂), characteristic signal at δ 3.6 ppm (s, 1H, CH), & δ 5.93 ppm (s, 1H, NH), δ 3.71 ppm (s, 2H, CH₂), δ 2.26 ppm (s, 3H, CH₃), δ 1.91 ppm (s, 3H, CH₃ of acetyl group), δ 6.03 ppm (s, 1H, =CH)[**14,15**]. The structure (**7b**) was consider most likely based on mass spectrum resulted in Mass spectra of (**7b**) reveals a molecular ion [M] + peaks at $m/z = 928$, and base peak at $m/z = 517$ [**16**]

3.7. SPECTRAL BEHAVIOUR

Spectral behaviours of the all syn the sized cyanine dyes in 95% ethanol were recorded using 1 cm cells. A stock solution (10⁻³ M) of the dyes was diluted to appropriate volume in order to obtain the required concentrations

REFERENCES

- [1] L Kong , Y., Liu, H., Wang, X he Tian, Q., Chen, T. Yu Peng, Li, Sheng, M. J. S. Dewar & N., Trinajstic, *J. Chem. Soc. A*, 1754, (1969).
- [2] E. Ravindran, and S. J., Ananthakrishnan, *J. Mater. Chem. C*, 3, 4359 (2015),
- [3] M . D., Bonifazi, *Chem. Soc.Rev.*, 41, 211. (2012).
- [4] O., Fenwick, C. V. K. Dyck, D., Murugavel, F. R. Cornil, & S., J. Haar, *Mater. Chem. C*, 3, 3007. (2015).
- [5a] Li., Yongjun, Liu, Taifeng, Liu, T Huibiao, Mao-Zhong,. & Li.; Yuliang, *Acc. Chem. Res.*, 47 (4), 1186-1198, (2014), [b] Li, Yongjun, Liu, Taifeng Huibiao

- Liu, Mao-Zhong Tian, & Li Acc. Yuliang ,Xue Zhao-ming, & Ij-,Xian Yang, *J. Mater. Chem. C*, **4**, 2990
(2016), [6] K., Matsumoto, H., Katsura, J., Yamauchi, T. Uchida, K. Aoyama, and T., Machiguchi, *Bull. Soc. Chim. Fr*, **133**, 891 **(1996)**.
- [7] C. W.,Bird, *Tetrahedron*, **54**, 10179. (1998), *Chem. Res.* **47** (4), 1186-1198 **(2014)**,
- [8a] F. S., Kim , G. Q. Ren & S. A., Jenekhe, *Chem. Mater.* **23**, 682. **(2011)**.
- [8b] MK, Mohanty, R, Sridhar, & S.Y. Padmanavan, *Indian J.Chem***158**:1146 **(1977)**.
- [9] A. I., Koraiem, A. M. El-Shafei, & I. M. Abdellah, *Organic Chemistry an Indian journal*,**14**, (2), **(2018)**.
- [10] A. I., Koraiem, A. M. El-Shafei, & I. M., Abdellah, *international journal of advanced research in Science, Engineering & technology*, **5**, (5), **(2018)**.
- [11a] A. I., Koraiem, A. M. El-Shafei, & I. M., Abdellah, *international Journal of Organic Chemistry in press*, **8** **(2018)**,
- [11a] A.I. M Koraiem, H. A., Shindy, R M. Abu-El-Hamd, & M. A Ibrahim,. *Journal of Applicable Chemistry, Accepted in* **(2018)**.
- [12] M. M. Gommaa, *Ph.D. Thesis, Faculty of Science, Aswan University* **(2001)**.
- [13] L. J Bellamy, *The infrared spectra of complex molecules*, London; Methuen, **(1962)**.
- [14] F. Scheinman, *Nuclear magnetic resonance of complex Molecules*, Braunschweig: Vieweg and Sohn GmbH, vol.1. **(1970)**.
- [15] T. J.; Batterham, *¹HNMR spectra of simple heterocycles*” Wiley New York, **(1973)**.
- [16] Q. N., Porter, and J;Baldas, *“Mass Spectrometry of Heterocyclic Compounds “* Wiely, New York, **(1971)**

Table (1): Characterisation data for (2a-d & 3a-d).

Comp. No.	Nature of product				Mol. Formula (M. wt)	Analytical analysis (%) Calcd (found)		
	Heterocyclic Substituent A(B)	m.p. °C	Yield %	Colour		C	H	N
2a	pyridin-4-ium (quinolin-1-ium)	290	65	Brown	C ₂₉ H ₂₅ N ₃ O ₃ I ₂ 464	75.14 75.35	5.44 5.70	9.07 9.35
2b	quinolin-4-ium (quinolin-1-ium)	175	75	Black	C ₃₃ H ₂₇ N ₃ O ₃ I ₂ 514	77.17 77.35	5.30 5.60	8.18 8.45
2c	quinolin-1-ium (quinolin-1-ium)	255	65	Buff	C ₃₃ H ₂₇ N ₃ O ₃ I ₂ 514	77.17 77.45	5.30 5.56	8.18 8.45
2d	quinolin-4-ium (pyridin-1-ium)	195	70	Brownish Red	C ₂₉ H ₂₅ I ₂ N ₃ O ₃ 717	48.56 48.66	3.51 3.67	5.86 5.99
3a	pyridin-4-ium (quinolin-1-ium)	> 300	75	Dark Brown	C ₃₄ H ₃₁ N ₃ O ₅ I ₂ 562	72.71 72.87	5.56 5.75	7.48 7.76
3b	quinolin-4-ium (quinolin-1-ium)	195	70	Black	C ₃₈ H ₃₃ N ₃ O ₅ I ₂ 612	74.62 74.87	5.44 5.65	6.87 6.99
3c	quinolin-1-ium (quinolin-1-ium)	> 300	80	Dark brown	C ₃₈ H ₃₃ N ₃ O ₅ I ₂ 612	74.62 74.78	5.44 5.65	6.87 6.99
3d	quinolin-4-ium (pyridin-1-ium)	180	90	Brownish red	C ₃₄ H ₃₁ I ₂ N ₃ O ₅ 815	50.08 50.35	3.83 4.00	5.15 5.35

Table (2): Characterisation data for (4a-d,5a-d):

Comp. No.	Nature of Product				Mol. Formula (M. wt)	Analytical Analysis (%) Calcd (found)			Absorption spectra In 95 % EtOH	
	Heterocyclic Substituent A(B)	m.p. °C	Yield %	Colour		C	H	N	λ_{\max} n. m	ϵ_{\max} mol ⁻¹ cm ²
4a	A=pyridin-4-ium- (quinolin-1-ium)	230	65	Brownish red	C ₃₃ H ₂₅ N ₃ O ₃ I ₂ 512	77.48 77.75	4.93 5.05	8.21 8.45	455 370	2000 4000
4b	A=quinolin-4-ium- (quinolin-1-ium)	285	70	Deep red	C ₃₇ H ₂₇ N ₃ O ₃ I ₂ 562	79.13 79.35	4.85 4.97	7.48 7.65	465 375	8800 8400
4c	A=quinolin-1-ium- (quinolin-1-ium)	195	55	Deep red	C ₃₇ H ₂₇ N ₃ O ₃ I ₂ 562	79.13 79.45	4.85 4.98	7.48 7.67	460 373	4000 7400
4d	A=quinolin-4-ium- (pyridin-1-ium)	270	65	Brownish red	C ₃₃ H ₂₅ N ₃ O ₃ I ₂ 512	77.48 77.75	4.93 5.08	8.21 8.45	475	13200
5a	A=pyridin-4-ium- (quinolin-1-ium)	280	75	Dark brown	C ₃₁ H ₂₃ I ₂ N ₃ O ₂ 723	51.47 51.67	3.21 3.45	5.81 5.94	510 445 415	(6000) (8000) (9600)
5b	A=quinolin-4-ium- (quinolin-1-ium)	>300	80	Deep Violet	C ₃₅ H ₂₅ N ₃ O ₂ I ₂ 520	80.91 81.00	4.85 5.05	8.09 8.35	660 570 470 435	(2400) (5400) (7800) (8800)
5c	A=quinolin-1-ium- (quinolin-1-ium)	290	70	Brown	C ₃₅ H ₂₅ N ₃ O ₂ I ₂ 520	80.91 81.00	4.85 5.08	8.09 8.34	512 450	(6800) (14000)
5d	A=quinolin-4-ium- (pyridin-1-ium)	>300	65	Dark Blue	C ₃₁ H ₂₃ I ₂ N ₃ O ₂ 723	51.47 51.67	3.21 3.55	5.81 5.99	690 595 560 400	(3200) (12800) (11400) (9600)

Table (3): Characterisation data (6a-f & 7a-f)

Comp. No.	Nature of Product				Mol. Formula (M.wt)	Analytical Analysis % Calcd (Found)			Absorption Spectra in 95% EtOH		
	Heterocyclic Substituent A[A' (B)]	M.P °C	Yield %	Colour		C	H	N	λ_{\max} n.m	ϵ_{\max} mol ⁻¹ cm ²	Absorption Range λ_{\max} n.m
6a	pyridin-4-ium[quinolin-4-ium(quinolin-1-ium)]	280	65	Deep red	C ₄₀ H ₃₂ I ₂ N ₄ O ₃ 870	55.19 55.54	3.71 3.86	6.44 6.67	610 380	(1800) (9800)	380-610
6b	quinolin-4-ium[quinolin-4-ium(quinolin-1-ium)]	230	63	Violet	C ₄₈ H ₃₆ N ₄ O ₃ I ₂ 716	80.43 80.67	5.06 5.45	7.82 7.98	690 575 460	(2000) (6000) (4200)	460-690
6c	quinolin-1-ium[quinolin-4-ium(quinolin-1-ium)]	235	75	Brownish red	C ₄₈ H ₃₆ N ₄ O ₃ I ₂ 716	80.43 80.65	5.06 5.56	7.82 7.98	635 ^{sh} 465 390	(2500) (2000) (7000)	390-635 ^{sh}
6d	quinolin-4-ium,[quinolin-4-ium(pyridin-1-ium)]	260	70	Deep violet	C ₄₄ H ₃₄ I ₂ N ₄ O ₃ 920	57.41 57.68	3.72 3.85	6.09 6.35	700 600 470	(2600) (4600) (10800)	470-700
6e	quinolin-4-ium[pyridin-4-ium(quinolin-1-ium)]	195	80	Brownish red	C ₄₄ H ₃₄ N ₄ O ₃ I ₂ 666	79.26 79.45	5.14 5.46	8.40 8.55	470 335	(3500) (5500)	335-470
6f	quinolin-4-ium[quinolin-1-ium(quinolin-1-ium)]	125	75	Shiny Red	C ₄₈ H ₃₆ N ₄ O ₃ I ₂ 716	80.43 80.67	5.06 5.45	7.82 7.99	500	(13000)	500
7a	pyridin-4-ium[quinolin-4-ium(quinolin-1-ium)]	105	65	Deep Red	C ₄₂ H ₃₂ I ₂ N ₄ O ₂ 878	57.42 57.55	3.67 3.83	6.38 6.56	620 390	(1000)	390-620
7b	quinolin-4-ium[quinolin-4-ium(quinolin-1-ium)]	235	60	Brownish red	C ₄₆ H ₃₄ I ₂ N ₄ O ₂ 674	81.88 81.98	5.08 5.35	8.30 8.45	690 585 470	(36000) (50000) (50000)	470-690
7c	quinolin-1-ium[quinolin-4-ium(quinolin-1-ium)]	230	60	Red	C ₄₆ H ₃₄ I ₂ N ₄ O ₂ 675	81.88 81.98	5.08 5.36	8.30 8.56	645 ^{sh} 475 400	(7000) (13500)	400-645 ^{sh}
7d	quinolin-4-ium,[quinolin-4-ium(pyridin-1-ium)]	220	65	Brownish red	C ₄₂ H ₃₂ N ₄ O ₂ I ₂ 625	80.75 80.87	5.16 5.35	8.97 9.05	700 620 490	(5000) (7500)	490-700
7e	quinolin-4-ium[pyridin-4-ium(quinolin-1-ium)]	250	55	Pink	C ₄₂ H ₃₂ N ₄ O ₂ I ₂ 625	80.75 80.89	5.16 5.45	8.97 9.05	480 355	(3500) (5000)	355-480
7f	quinolin-4-ium[quinolin-1-ium(quinolin-1-ium)]	160	80	Deep Red	C ₄₆ H ₃₄ I ₂ N ₄ O ₂ 675	81.88 81.98	5.08 5.25	8.30 8.55	520	(5500) (7900) (7900)	520

Table (4) : Comparison of Absorption Spectra in EtOH of (5a-d & 7a-f)

Comp. No.	Heterocyclic Substituent A(B)	Absorption Spectra of in 95% EtOH		
		λ_{\max} n.m	ϵ_{\max} mol ⁻¹ cm ²	Range λ_{\max} n.m
5a	A=pyridin-4-ium-(quinolin-1-ium)	510,445,415	(6000),(8000) (9600)	415-510
5b	A=quinolin-4-ium-(quinolin-1-ium)	660,570,470,435	(2400),(5400) (7800),(8800)	435-660
5c	A=quinolin-1-ium-(quinolin-1-ium)	512, 450	(6800),(14000)	450-512
5d	A=quinolin-4-ium-(pyridin-1-ium)	690,595,560,400	(3200),(12800) (11400),(9600)	400-690
7a	A=pyridin-4-ium[quinolin-4-ium(quinolin-1-ium)]	620 390	(1000)	390-620
7b	A=quinolin-4-ium[quinolin-4-ium(quinolin-1-ium)]	690,585 470	(36000), (50000) (50000)	470-690
7c	A=quinolin-1-ium[quinolin-4-ium(quinolin-1-ium)]	645 ^{sh} ,475 400	(7000),(13500)	400-645 ^{sh}
7d	A=quinolin-4-ium,[quinolin-4-ium(pyridin-1-ium)]	700,620 490	(5000), (7500)	490-700
7e	A=quinolin-4-ium[pyridin-4-ium(quinolin-1-ium)]	480,355	(3500), (5000)	355-480
7f	A=quinolin-4-ium[quinolin-1-ium(quinolin-1-ium)]	520	(5500), (7900) (7900)	520